arXiv:1211.4862v1 [quant-ph] 20 Nov 2012

Planar quantum squeezing via quantum non-demolition measurements in cold atomic ensembles

Graciana Puentes[1], **Giorgio Colangelo**[1], **Robert J. Sewell**[1], **and Morgan W. Mitchell**[1,2]

[1] ICFO - Institut de Ciencies Fotoniques, Mediterranean Technology Park, 08860 Castelldefels, Barcelona, Spain

[2] ICREA - Institucio Catalana de Recerca i Estudis Avancats, 08015 Barcelona, Spain

E-mail: `graciana.puentes@icfo.es`

Abstract. Planar quantum squeezed states, i.e. quantum states which are squeezed in two orthogonal components on a plane, have recently attracted attention due to their applications in atomic interferometry and quantum information [1, 2]. Unlike the quadratures of the radiation field which are inherently transverse, simultaneous squeezing in two orthogonal spin components can be achieved due to the fact that angular momentum belongs to SU(2,3) group, a feature which can be exploited to tailor the corresponding commutator between two such operators, and reduce orthogonal variances simultaneously. In this paper, we present a novel scheme for planar quantum squeezing via quantum non-demolition (QND) measurements in spin-1 systems [3]. The QND measurement is achieved via paramagnetic Faraday rotation on off-resonant probe light, and the planar quantum squeezing is obtained with the aid of an external magnetic field which introduces a Larmor precession on the plane where squeezing takes place. We show that planar quantum squeezed states can be used to reconstruct an arbitrary quantum parameter, such as a phase, without any prior information, and could therefore find relevant applications in single-shot atomic magnetometry where temporal fluctuation can prevent the implementation of iterative procedures.

1. Introduction

Quantum squeezing, the reduction of uncertainty in quantum observables below "classical" or standard quantum limits, is an area of active research with applications in quantum information processing, quantum communications, and quantum metrology. Here we consider a practical, measurement-based strategy to produce a new kind of squeezing in atomic spin ensembles. By way of introduction, we note that a spin **F** obeys the spin uncertainty relations $\Delta F_z \Delta F_x \geq \frac{1}{2}|\langle F_y \rangle|$ and permutations, which are distinctly different from the Heisenberg relation $\Delta X \Delta P \geq \frac{1}{2}$ that governs canonical variables X, P [4]. For canonical variables, the constant RHS of the Heisenberg relation implies that reduction of the variance in one quadrature inevitably increases the variance of the other. In contrast, in spin systems the RHS of the uncertainty relation may vanish, e.g. if $\langle F_y \rangle = 0$, with the consequence that two spin components, e.g. F_z and F_x, may be *simultaneously* squeezed, with the uncertainty being absorbed by the third component. An analogous situation involving the Stokes parameters of light, which themselves obey angular momentum commutation relations, has also been studied [5, 6]. For atomic spin ensembles, the suggestion of "intelligent" spin states [7] predates even the concept of optical squeezing, and "planar quantum squeezed states" (PQS) have been explored in depth [1, 2].



Figure 1. Illustration of spin variances for: a) an unsqueezed state with SQL variances $\Delta^2 F_i$ for all components i $\in \{x, y, z\}$, b), b') a state with squeezed azimuthal component. For measurements of Fz, this squeezing is advantageous only for certain precession angles, as in b), but disadvantageous for others, as in b'). c), c') a planar squeezed state with both variances $\Delta^2 F_x$ and $\Delta^2 F_z$ reduced below the SQL is advantageous at all angles.

As described in [1, 2], planar quantum squeezing (PQS) has the prospect of improving the precision of atomic interferometers at arbitrary phase angles, as it requires both orthogonal variances to be below the shot-noise level and with the same scaling. This is opposed to squeezing on a single direction, which can only be applied to determine a particular phase angle, known *a priori* (see Fig. 1). In addition, PQS states reduce phase noise everywhere on a plane in a single shot, and can therefore be employed in cases where iterative approaches are not applicable. Thus they have promising applications in high-bandwidth atomic magnetometry. In particular, PQS can also be of interest in situations when the phase is not known to a good approximation or when such phase does not remain constant during repeated measurements, as it requires a minimal number of measurements. Such measurement strategies are radically different from those employing multiple sequential measurements, which inherently assume that the phase-shift is a time-invariant, classical parameter.

He *et al.* [1] proposed an implementation of PQS using a two-well Bose condensate with tunable and attractive interactions. Here, we propose an alternative system which can provide PQS, and which has potential applications in high bandwidth atomic magnetometry on a plane, without the need for prior information, and which can be employed even in the presence of temporal fluctuations, where iterative approaches are not applicable. In this approach, the PQS state is prepared by stroboscopic probing a polarized ensemble of cold ^{87}Rb atoms, with the aid of a constant external magnetic field which provides the planar Larmor precession required in order to squeeze two orthogonal components by sequential measurements. We work with the extended collective angular momentum variable **F** for spin-1 systems. The collective atomic spin is measured using paramagnetic Faraday rotation with an off-resonant probe. The ensemble of spins, polarized along a fixed direction by optical pumping, interacts with an optical pulse of duration τ and polarization described by the Stokes vector $\mathbf{S} = (\hat{S}_x, \hat{S}_y, \hat{S}_z)$, through an effective Hamiltonian.

The article is structured as follows: First, in Section II, we present the basic formalism for light-atom interactions in polarized atomic ensembles. Second, in Section III, we report numerical evidence of PQS in our system. Next, in Section IV, we provide an example for the applications of these ideas in high bandwidth (single shot) atomic magnetometry. Finally, in Section V we present our conclusions.

2. Atom-light Interaction

Our modeling is based on covariance matrix techniques introduced by Madsen and Mølmer [8, 9], extended to include inhomogeneities by Koschorreck *et al.* [10], and to spin-1 atoms by Colangelo *et al.* [3]. In brief, we consider a magnetic field $\mathbf{B}$ of unknown magnitude oriented along the y axis, in interaction with a spin-1 ensemble, e.g. ^{87}Rb in the $F = 1$ ground state, and probed using Faraday rotation probing by near-resonant light pulses. The ensemble is described by collective spin observables, for example $\hat{F}_x \equiv \sum_i \hat{f}_x^{(i)}$, where $\hat{f}_x^{(i)}$ indicates the x-component of the spin of the i'th atom. We include in the analysis both spin orientation $\hat{F}_x, \hat{F}_y, \hat{F}_z$, and also other spin variables $\hat{J}_x, \hat{J}_y, \hat{J}_k, \hat{J}_l, \hat{J}_m$ defined in terms of $\hat{f}_y$, $\hat{f}_z$, $\hat{j}_x \equiv \hat{f}_x^2 - \hat{f}_y^2$, $\hat{j}_y \equiv \hat{f}_x\hat{f}_y + \hat{f}_y\hat{f}_x$, $\hat{j}_k \equiv \hat{j}_k = \hat{f}_x\hat{f}_z + \hat{f}_z\hat{f}_x \quad \hat{j}_l = \hat{f}_y\hat{f}_z + \hat{f}_z\hat{f}_y \quad \hat{j}_m = \frac{1}{\sqrt{3}}(2\hat{f}_z^2 - \hat{f}_x^2 - \hat{f}_y^2)$, respectively. The optical pulses are described in terms of Stokes operator components $\hat{S}_x, \hat{S}_y, \hat{S}_z$, defined as:

$$\hat{S}_x \equiv \frac{\hbar}{2}\hat{\mathbf{a}}^\dagger\sigma_x\hat{\mathbf{a}} \quad \hat{S}_y \equiv \frac{\hbar}{2}\hat{\mathbf{a}}^\dagger\sigma_y\hat{\mathbf{a}} \quad \hat{S}_z \equiv \frac{\hbar}{2}\hat{\mathbf{a}}^\dagger\sigma_z\hat{\mathbf{a}}. \tag{1}$$

Where $\hat{\mathbf{a}} \equiv (\hat{a}_+, \hat{a}_-)^T$ and $\hat{a}_+, \hat{a}_-$ are the annihilation operators for the left and right circular polarization and σ_x, σ_y, σ_z the Pauli matrices. The $\hat{S}_x$, $\hat{S}_y$ and $\hat{S}_z$ Stokes operators represent polarized light in the horizontal or vertical direction, polarized light in the $\pm 45^o$ direction, and right hand or left hand circularly polarized light, respectively. As mentioned, they obey the same commutation relations as angular momentum operators.

We compute the evolution of the phase-space vector

$$\hat{\mathbf{v}} = [\hat{F}_x, \hat{F}_y, \ldots, \hat{J}_m, \hat{S}_x, \hat{S}_y, \hat{S}_z] \tag{2}$$

and the covariance matrix

$$\gamma = \frac{1}{2}\langle \hat{\mathbf{v}} \otimes \hat{\mathbf{v}} + (\hat{\mathbf{v}} \otimes \hat{\mathbf{v}})^{\mathbf{T}} \rangle - \langle \hat{\mathbf{v}} \rangle \otimes \langle \hat{\mathbf{v}} \rangle, \tag{3}$$

which together provide a full description of the state. The components of the phase-space vector and covariance matrix have initial values computable from the initial atom-light state, typically given by a collective atomic spin fully polarized along the direction of the $\hat{F}_x$ component, and probe light linearly polarized along the Stokes vector component $\hat{S}_x$. The phase-space vector evolves as

$$\hat{\mathbf{v}}(t + \tau) = \mathbf{T}_\tau \hat{\mathbf{v}}(t) \tag{4}$$

where $\mathbf{T}_\tau$ accounts for coherent evolution under the interaction Hamiltonian

$$\tau H_{\text{eff}} = \tau\mu_0 B_y \cdot \hat{F}_y + G_1\hat{S}_z\hat{F}_z + G_2(\hat{S}_x\hat{J}_x + \hat{S}_y\hat{J}_y) \tag{5}$$

and loss of amplitude (damping) due to decoherence and dephasing, caused by incoherent scattering of probe photons and inhomogeneous magnetic fields, respectively [10]. The covariance matrix evolves as

$$\gamma(t+\tau) = \mathbf{T}_\tau \gamma(t) \mathbf{T}_\tau^{\mathbf{T}} + N_\tau . \tag{6}$$

where N_τ is a noise matrix which describes the noise associated with damping processes [10, 3], and we have taken $\hbar = 1$ for simplicity.

The details of the covariance matrix formalism for quantum non-demolition (QND) measurements in cold atomic ensembles for spin $F = 1$ systems, in the presence of an external $\mathbf{B}$-field, are detailed in Ref. [3]. Here we limit to mention that in the presence of a dynamical decoupling scheme it is possible to eliminate the tensorial G_2 component in H_{eff} and achieve a truly QND measurement via Faraday rotation of the probing light. In this case, by reducing the variance in the light pulses via a precise measurement of the Stokes operator $\hat{\mathbf{S}}_{\mathbf{y}}$ it is possible to squeeze (or reduce the variance) on the spin atomic component $\hat{\mathbf{F}}_{\mathbf{z}}$ in a QND fashion, due to the vectorial coupling term G_1 in $\hat{H}_{\text{eff}}$. In addition, by measuring at time intervals given by a quarter of the Larmor period, it is possible to squeeze the orthogonal atomic component $\hat{\mathbf{F}}_{\mathbf{x}}$, due to the external magnetic field B_y which introduces a Larmor precession on the (x, z)-pane. In the next section, we present numerical results based on such formalism for the realization of planar quantum squeezing in the system.

3. Numerical Results

In this Section, we present numerical results validating the use of QND measurements in cold atomic ensembles for conditional preparation of planar quantum squeezed states. The operational definition we adopt for planar quantum squeezing follows the approach by He *et al.* [1, 2], and can be summarized as follows. We require that the normalized sum of the variances on a plane should be below the Standard Quantum Limit (SQL). For the case of spin-1 ($F = 1$) systems, this reads:

$$0 < \frac{\Delta F_{||}}{\langle |F| \rangle} \equiv \frac{\sqrt{\Delta F_x^2 + \Delta F_z^2}}{\langle |F| \rangle} < 1. \tag{7}$$

In addition to this, we require that the individual uncertainties should be below the SQL $\Delta F_x < \langle |F| \rangle$, and $\Delta F_z < \langle |F| \rangle$. In order to achieve such kind of state, the angular momentum component in the direction perpendicular to the plane should be zero ($\langle F_y \rangle = 0$) so that the uncertainties on the plane are uncorrelated $\Delta F_x \Delta F_z \geq 0$. Furthermore, we require the mean value of the angular momentum on the plane to be large so that we can reduce the uncertainty in two components, at the expenses of increasing the noise in the third component, i.e., $\Delta F_{z(x)} \Delta F_y \geq \frac{\langle F_{x(z)} \rangle}{2}$.

We performed numerical simulations for stroboscopic probing [3],[10], in a cold atomic ensemble containing $N_{at} = 10^6$ ^{87}Rb atoms with coupling parameter $G_1 = 0.5 \times 10^{-7}$ corresponding to a detuning in the off-resonant probe given by $\Delta = -2\pi \times 2$ GHz. The atomic ensemble is initially polarized in the x-direction with 99.9 % Fidelity, and technical noise due to atom number fluctuations given by $\Delta N = 0.0007$. The initial collective atomic spin mean values are $\langle F_x \rangle_{t=0} = N_{at}$, $\langle F_y \rangle_{t=0} = 0$, and $\langle F_z \rangle_{t=0} = 0$. Additionally, we introduced an external magnetic field $B_y = 20.042$ mG, which sets a Larmor precession in the (x, z)-plane, with a characteristic period of ${}_L \approx 71$ μs. We chose this magnitude for B_y as an upper limit, in order to allow sufficient time within a Larmor cycle for stroboscopic probing. The experimental scheme is illustrated in Fig. 2, and was described in detail in Ref. [14]. The atomic ensemble was probed at four locations during the Larmor cycle by a single off-resonant optical pulse at each location. The atomic cloud was left free to precess at time intervals given by $T_L/4$ in order to probe the collective atomic spin sequentially on a plane $(F_z, F_x, -F_z, -F_x)$, as required for planar quantum squeezing. The length of optical pulses in the sequences was $\tau = 1\mu$s, and each pulse contained 0.5×10^8 photons. The optimal number of photons on each individual pulse was obtained as a trade-off between maximal squeezing and minimum loss of atomic polarization due to optical scattering. We note that in article we did not consider a dynamical decoupling scheme, though further squeezing could in principle be achieved by using a dynamical decoupling approach [12].



Figure 2. Experimental scheme for planar quantum squeezing via quantum non-demolition measurements. The atomic ensemble is initially polarized along the x-direction, and precesses in the (x, z)-plane due to an external field B_y which introduces a Larmor rotation. Planar quantum squeezing is obtained by measuring the Faraday rotation of the Stokes components in the off-resonant probe light at four different spatial locations during the Larmor precession cycle [14].

The numerical results are shown in Fig. 3 to Fig. 5. In Figure 3, black curve corresponds to $\langle F_x \rangle$, blue curve corresponds to $\langle F_z \rangle$, and red curve is $\langle F_y \rangle$. The ensemble of atoms is initially polarized in x-direction, and precesses in the (x, z)-plane due to the B-Field at the Larmor frequency. The mean value in the direction perpendicular to the plane (red curve) is always approximately zero, i.e., $\langle F_y \rangle \approx 0$, as required for planar quantum squeezing (i.e. $\Delta F_x \Delta F_z \geq 0$). This allows for the variances in the (x, z) plane to be independently reduced. The deviations from $\langle F_y \rangle = 0$ are due to the tensorial coupling term in the effective Hamiltonian $G_2(\hat{S}_x \hat{J}_x + \hat{S}_y \hat{J}_y)$, and could be removed via dynamical decoupling [12].

Next, we evaluated the normalized variances of the collective angular spin components after sequential stroboscopic probing on a plane. This is shown in Fig. 4. We note that the individual normalized variances on the (x, z)-plane, corresponding to blue curve and black curve are initially equal to unity due quantum and technical noise. They oscillate as a consequence of the Larmor precession, and are sequentially reduced due to the stroboscopic probing, as indicated by the discontinuous jumps in $\Delta^2 F_y$ (red curve). On the other hand, the variance in the orthogonal direction $\Delta^2 F_y$ is sequentially increased well above the SQL (dashed cyan line), as expected for a planar quantum squeezed (PQS) state. The ideal planar quantum squeezed state, for the system parameters chosen, is achieved at approximately 500 μs.

Figure 5 shows the normalized planar squeezing parameter $\frac{\Delta F_{||}}{\langle |F| \rangle}$. It is seen that for times longer than 200μs, this parameter is always bellow the unit SQL, and is sequentially reduced to less than $\frac{\Delta F_{||}}{\langle |F| \rangle} < 0.8$, with each individual components squeezed to $\frac{\Delta F_x}{\langle |F| \rangle} \approx \frac{\Delta F_y}{\langle |F| \rangle} \approx 0.5$ which con-



Figure 3. Normalized mean values of collective atomic angular momenta. Blue curve $\langle F_z \rangle$, black curve $\langle F_x \rangle$, and red curve $\langle F_y \rangle$. The mean value in the direction orthogonal to the plane of precession remains always approximately to zero $\langle F_y \rangle \approx 0$, as required for planar quantum squeezing.



Figure 4. Normalized variance of collective atomic spins. Blue curve $\frac{\Delta F_z}{\langle |F| \rangle}$, black curve $\frac{\Delta F_x}{\langle |F| \rangle}$, and red curve $\frac{\Delta F_y}{\langle |F| \rangle}$. At all times the variances on the (x, z)-plane are below the standard quantum limit (SQL=1) indicated by the dashed line, and are decreased at the expense of increasing noise in the orthogonal direction (red curve). The discontinuities in ΔF_y correspond to stroboscopic probings. Maximal planar quantum squeezing is achieved at approximately 500μs.

firms the presence of planar quantum squeezing in the system. Inset in Fig. 5 shows a reduction by 40% in the normalization factor $|F|$ due to optical scattering during the probing sequence, which in turns limits the maximal planar squeezing that can be obtained.

3.1. Limits on planar quantum squeezing due to optical scattering

In the absence of optical scattering and spontaneous emission the squeezing that can be obtained by QND measurements is of order $\xi = 1/(1 + \kappa)$, where $\kappa \propto G_1^2$ is the atom-light coupling parameter. The quantum uncertainty after squeezing can be expressed as $\Delta F = \xi \times N_{at}/2$, where



Figure 5. Normalized planar squeezing parameter $\frac{\Delta F_{||}}{\langle |F| \rangle}$. For times longer than 200 μs the planar uncertainty remains below the standard quantum limit (SQL=1), as indicated by the dashed line, and is reduced sequentially due to stroboscopic probing. Maximal planar quantum squeezing is achieved at approximately 500 μs and is reduced at longer times due to loss of atomic polarization by optical scattering. Inset shows the normalization factor $|F|$, which is reduced by 40 % during the total measurement time.

$N_{at}/2$ is the nominal uncertainty for a coherent state in a spin-1 system. This would suggest that infinite squeezing can in principle be achieved from Kerr interaction. However, there are severe limitations on the amount of squeezing that can be achieved due to spontaneous emission. In fact, the coupling parameter κ can also be expressed as $\kappa = \alpha\eta$, where α is the atomic ensemble optical column density, and $\eta << 1$ is the spontaneous emission rate. A crude estimate for the squeezing parameter in the presence of optical scattering can be expressed by $\xi = 1/(1+\alpha\eta)+2\eta$, and it is minimum for an optimal spontaneous decay probability $\eta_0 = 1/\sqrt{2\alpha}$. For a typical system with optical density $\alpha \approx 25$ this optimal value gives an lower bound on the amount of squeezing given by roughly $\xi^{\mathrm{min}} \approx 0.5$ (3dB of noise reduction) [15].

A similar argument can be employed to determine a lower bound on the squeezing parameter that can be achieved for planar quantum squeezing via QND measurements. Following the definition of $\Delta F_{||} = \sqrt{\Delta F_x^2 + \Delta F_z^2}$, we can define a similar expression for the planar squeezing parameter $\xi_{||}$ in terms of the individual squeezing parameters $\xi_{x,z}$, in the following form:

$$\xi_{||} = \sqrt{\xi_x^2 + \xi_z^2}, \tag{8}$$

where $\xi_x = 1/(1+\alpha_x\eta)+2\eta$ and $\xi_z = 1/(1+\alpha_z\eta)+2\eta$. Assuming the same optical depth in both directions $\alpha_x = \alpha_z = 25$, we can estimate the minimum planar squeezing parameter to be $\xi_{||}^{\mathrm{min}} \approx 0.711$ (roughly 2 dB of noise reduction). In general, the planar squeezing parameter results larger than the squeezing parameter in a single direction by roughly a factor of $\sqrt{2}$ for the case of symmetrical optical column densities. If the optical depths in each direction in the plane are different, the planar squeezing parameter will change. In particular, in order to achieve $\xi_{||} < 1$ it is required sufficient a optical depth in both directions, and both α_x and α_y should be above 10 for the case of symmetric densities. This is depicted in Figure 6, for experimentally feasible values for the optical depths $0 < \alpha_{x,z} \leq 100$.



Figure 6. Planar squeezing parameter $\xi_{||}$ in terms of optical depths ($\alpha_{x,z}$). Both α_x and α_z should be above 10 in order to achieve a planar squeezing parameter below the standard quantum limit $\xi_{||} < 1$.

3.2. Optimal measurement strategy in the presence of noise

The optimal measurement strategy will depend in general on the amount of initial quantum and technical noise in the angular momentum components $\mathrm{var}(F_i(t=0))$, with $i = (x, z)$. If the initial noise in one component is sufficiently low (i.e., below the SQL), then it seems enough to squeeze the remaining component by probing in a single direction. However, in a more realistic scenario the initial noise can be arbitrarily large in either component, such as for a coherent state, in which case it is required to probe two orthogonal directions on a plane in order to obtain a planar quantum squeezing parameter below the SQL ($\Delta F_{||}/\langle|F|\rangle < 1$), such situation corresponds to actual PQS. This is clearly shown in Figure 7. We performed simulations considering initial quantum noise corresponding to a polarized initial state with different Fidelities, and different amounts of initial random technical noise. Black curve corresponds to $\Delta N \approx 0.0001$, blue curve corresponds to $\Delta N \approx 0.0004$, red curve corresponds to $\Delta N \approx 0.0007$, and green curve $\Delta N \approx 0.001$. The inset depicts the measurement strategy, given by probing a single component ($F_z, -F_z$) at interval times given by half of the Larmor period ($T_L/2$) (dashed lines), or probing two orthogonal components ($F_z, F_x, -F_z, -F_x$) at interval times given by a quarter of such period ($T_L/4$) (solid lines). As expected, in the presence of initial quantum and technical noise above the SQL (blue, red and green curves), larger planar quantum squeezing can be achieved by probing the system sequencially on a plane.

4. Applications: Atomic magnetometry

Experimentally determined phase-shifts are never truly classical, nor time-invariant [2]. Phases often vary with time, and can not always avoid measurement back action, a limitation that can rule out the possibility of iterative techniques. The physical reason for this is that phase-shifts are related to energy-shifts terms in the radiation field Hamiltonian [16], a feature which has dynamical consequences.

In this Section, we describe a particular application of planar quantum squeezing (PQS) in



Figure 7. Normalized planar quantum squeezing parameter $\Delta F_{||}/\langle|F|\rangle$ for different amounts of initial quantum and technical noise, and for two different measurement strategies. Dashed lines corresponds to stroboscopic probing on a single direction $(F_z, -F_z)$, and solid lines corresponds to stroboscopic measurement on a plane $(F_z, F_x, -F_z, -F_x)$, as illustrated at the inset. Black, blue, red, and green curves correspond to $\Delta N = 0.0001$, $\Delta N = 0.0004$, $\Delta N = 0.0007$, and $\Delta N = 0.001$. As expected, for a sufficient amount of initial noise (blue, red and green curves) a larger amount of planar quantum squeezing ($\Delta F_{||}/\langle|F|\rangle < 1$) can be achieved by stroboscopic probing on a plane (solid lines).

atomic magnetometry, for determination of *arbitrary* quantum phase parameters below the SQL, without the need for prior information or without the requirement of linearized fluctuations. PQS states can be used to determine an almost arbitrary phase angle ϕ, within a finite interval along the precession of the collective atomic magnetic moment $\mathbf{F}$. The precession of this vector is determined by a phase $\phi = \omega_L t$, where ω_L is the Larmor frequency, which in turn is proportional to the external magnetic field B_{ext} applied on the system, so a high precision in the estimation of parameter ϕ can be mapped onto precision magnetometry. For an external field along the y-direction B_y, we expect a precession of the collective atomic spin on the $(\hat{x}, \hat{z})$ plane. The rotation of the collective spin on the plane can be written as:

$$F(\phi) = F_x \cos(\phi) + F_z \sin(\phi). \tag{9}$$

We can calculate the parameter uncertainty via $\Delta\phi = \frac{\sqrt{\Delta^2 F(\phi)}}{|dF(\phi)/d\phi|}$. By differentiating $F(\phi)$ we obtain:

$$\Delta\phi = \frac{\sqrt{\Delta^2 F(\phi)}}{|F_z \cos(\phi) - F_x \sin(\phi)|}, \tag{10}$$

where $\Delta^2 F(\phi) = \Delta^2 F_x \cos(\phi)^2 + \Delta^2 F_z \sin(\phi)^2 + [\text{cov}(F_x, F_z) + \text{cov}(F_z, F_x)] \sin(\phi)\cos(\phi)$. Here $\text{cov}(F_i, F_j)$ stands for the covariance of the operators, i.e. $\langle F_i F_j \rangle - \langle F_i \rangle \langle F_j \rangle$. If we work in the basis where $\text{cov}(F_x, F_z) = \text{cov}(F_z, F_x) = 0$ (this is in principle always possible), we can write $\Delta^2 F(\phi) = \Delta^2 F_x \cos(\phi)^2 + \Delta^2 F_z \sin(\phi)^2$. We can now set a bound on $\Delta^2 F(\phi)$ in terms of planar squeezing such that:

$$\Delta^2 F(\phi) = \Delta^2 F_x \cos(\phi)^2 + \Delta^2 F_z \sin(\phi)^2 \leq \Delta^2 F_{||}, \tag{11}$$



Figure 8. Parameter uncertainty ($\Delta\phi$) below the standard quantum limit $\Delta\phi < 1/\sqrt{N} = 0.001$ for planar quantum squeezed state with different levels of initial quantum noise. Fig. 8 (a) considering $F_x = F_z$, and Fig 8 (b) considering $F_z = 0$. Black, blue, red, magenta and green curves correspond to planar squeezed variances given by $\Delta^2 F = 0.01 \times N_{at}/2$, $\Delta^2 F = 0.1 \times N_{at}/2$, $\Delta^2 F = 0.25 \times N_{at}/2$, $\Delta^2 F = 0.1 \times N_{at}/2$, and $\Delta^2 F = 0.5 \times N_{at}/2$, respectively. For a realistic PQS state obtained via QND measurement, with $\Delta^2 F = 0.25 \times N_{at}/4$ (magenta curve), it is possible to reconstruct an arbitrary parameter in the interval $\phi \in [-\pi/4, 0.6) \cup (1.2, \pi]$ for case (a), or an arbitrary parameter in the interval $\phi \in [-\pi/4, -0.2) \cup (0.3, \pi]$ for case (b).

therefore obtaining an upper bound to the quantum noise present in the magnetometer given by the planar variance $\Delta^2 F_{||}$, thus confirming that PQS states are optimal for quantum-phase estimation as they reduce the variance on a plane. We note that planar quantum squeezed states give the lowest planar variance for arbitrary angles on a finite interval, with the exception of the specific singular values which make the denominator in Eq. (10) equal to zero. This feature is depicted in Fig. 8 and Fig. 9.

Figure 8 shows the phase parameter uncertainty $\Delta\phi$ below the SQL ($\Delta\phi < 1/\sqrt{N} = 0.001$) for a PQS state as given in Eq. 10, for different values of the collective atomic spin components $F_{x,z}$, respectively. Fig 8 (a) corresponds to $F_x = F_y$ and Fig. 8 (b) corresponds to $F_z = 0$. Black, blue, red, magenta and green curves correspond to planar squeezed variances given by $\Delta^2 F = 0.01 \times N_{at}/2$, $\Delta^2 F = 0.1 \times N_{at}/2$, $\Delta^2 F = 0.25 \times N_{at}/2$, and $\Delta^2 F = 0.5 \times N_{at}/2$, respectively. For a realistic PQS state obtained via QND measurements with $\Delta^2 F = 0.25 \times N_{at}/2$ (magenta curve), it is possible to reconstruct an arbitrary parameter in the interval (a) $\phi \in [-\pi/4, 0.6) \cup (1.2, \pi]$, (b) $\phi \in [-\pi/4, -0.2) \cup (0.3, \pi]$, with precision below



Figure 9. Comparison of parameter uncertainty ($\Delta\phi$) below SQL $\Delta\phi < 1/\sqrt{N} = 0.001$ for a planar quantum squeezed state (solid line), squeezed state in a single component (dashed line), and coherent state (dashed-dotted line). Fig. 9 (a) considering atomic collective spins $F_x = F_z$, and (b) considering $F_z = 0$. For the planar quantum squeezed state the variances are given by $\Delta^2 F_x = \Delta^2 F_z = 0.25 \times N_{at}/2$, for the state squeezed on a single component $\Delta^2 F_x = 0.25 \times N_{at}/2$ and $\Delta F_z = N_{at}/2$, and for the coherent state we have $\Delta^2 F_x = \Delta^2 F_z = N_{at}/2$. (a) Planar quantum squeezed states allow for parameter estimation below the SQL at larger intervals than coherent states, or states squeezed on a single component. (b) For the case $F_x = 0$ or $F_z = 0$ the estimation interval for a planar quantum squeezed state is roughly comparable (and slightly higher) to the one given by a state squeezed by the same amount on a single component, as expected. This shows that for a given amount of squeezing, planar quantum squeezed states are optimal for parameter reconstruction at finite intervals.

the SQL.

Next, Figure 9 shows a comparison of the phase parameter uncertainty $\Delta\phi$ below the SQL ($\Delta\phi < 1/\sqrt{N} = 0.001$) for a PQS state (solid line), a squeezed state on a single component (dashed line), and a coherent state (dashed-dotted line), for different values of the collective atomic spin components $F_{x,z}$, respectively. Fig. 9 (a) corresponds to $F_x = F_z$, and Fig. 9 (b) corresponds to $F_z = 0$. For the planar quantum squeezed state the variances are given by $\Delta^2 F_x = \Delta^2 F_z = 0.25 \times N_{at}/2$, for the state squeezed on a single component $\Delta^2 F_x = 0.25 \times N_{at}/2$ and $\Delta F_z = N_{at}/2$, and for the coherent state we set $\Delta^2 F_x = \Delta^2 F_z = N_{at}/2$. In Fig. 9 (a) we see that planar quantum squeezed states (solid line) allow for parameter estimation below the SQL at broader intervals than coherent states, or states squeezed on a single component. Fig.

9 (b), in the case $F_x = 0$ or $F_z = 0$ the estimation interval for a planar quantum squeezed state is roughly comparable (and slightly higher) than the one given by a state squeezed by the same amount but on a single component, as expected, for a non-vanishing squeezed component. This shows that for a given amount of squeezing, planar quantum squeezed states are optimal for reconstructions of arbitrary parameters with no *a priori* information.

5. Conclusions

To conclude, we have demonstrated that a novel kind of planar quantum squeezed (PQS) state, which squeezes the variances on two orthogonal spin components, can be prepared in cold atomic ensembles, via quantum non-demolition (QND) measurements. This is possible due to the fact that angular momentum belongs to SU(2,3) group which contains at least three orthogonal spin components, unlike EM-field quadratures which are inherently transverse, and whose commutator is a constant [4]. Such PQS states have important applications in the determination of arbitrary quantum phases below the standard quantum limit (SQL), in a single shot, and are therefore promising candidates for high-bandwidth atomic magnetometry in cases where iterative approaches are not applicable due to dynamical fluctuations. Experimental demonstrations of such atomic PQS states are currently under progress.

ACKNOWLEDGEMENTS

This work was supported by the Spanish MINECO under the project MAGO (ref. no. FIS2011-23520) and by the European Research Council under the project AQUMET. G.P. acknowledges financial support from Marie-Curie International Fellowship COFUND.

[1] Q. Y. He, S-G. Peng, P. D. Drummond, and M. D. Reid, Phys. Rev. A **84**, 022107 (2011).
[2] Q. Y. He *et al*, N. Jour. Phys. **14**, 093012 (2012).
[3] G. Colangelo *et al*, in preparation for submission to New. J. Physics (2012).
[4] R. Loudon and P. L. Knight, J. Mod. Opt. **34**, 709 (1987).
[5] N. Korolkova *et al*, Phys. Rev. A **65**, 052306 (2002).
[6] R. Schnabel *et al*, PhysRevA.67.012316 (2003).
[7] C. Aragone, G. Guerri, S. Salamo, and J. L. Tani, J. Phys. A **7**, L149 (1974).
[8] L. B. Madsen, and K. Molmer, Phys. Rev. A **70**, 052324 (2004).
[9] K. Molmer, and L. B. Madsen, Phys. Rev. A **70**, 052102 (2004).
[10] M. Koschorreck and M. Mitchell, J. Phys. B **42**, 195502 (2009).
[11] D. V. Kupriyanov, O. S. Mishina, I. M. Sokolov, B. Julsgaard, and E. Polzik, Phys. Rev. A **71**, 032348 (2005).
[12] M. Koschorreck *et al*, Phys. Rev. Lett. **105**, 093602 (2010).
[13] M. Koschorreck, M. Napolitano, B. Dubost, and M. Mitchell, Phys. Rev. Lett. **104**, 093602 (2010).
[14] M. Napolitano, M. Koschorreck, B. Dubost, N. Behbood, R. J. Sewell, M. W. Mitchell, Nature 471, 486-489 (2011).
[15] K. Hammerer, K. Molmer, E. S. Polzik, and J. I. Cirac, Phys. Rev. A **70**, 044304 (2004).
[16] P. D. Drummond, Phys. Rev. A **34**, 2718 (1989).
[17] C. M. Caves, Phys. Rev. D **23**, 1693 (1981).